

Double-peak spectral energy density of GRBs and the true identity of GRB 031203

Shlomo Dado¹ and Arnon Dar¹

ABSTRACT

A double-peak spectral-energy-density of γ -rays, similar to that observed in blazars, is expected in γ -ray bursts (GRBs) produced in supernova (SN) explosions. The second peak, which is formed by inverse Compton scattering of the ambient SN light by cosmic ray electrons accelerated by the jets from the SN explosion, has a much higher peak-energy than the first ordinary peak. However, in X-ray flashes (XRFs), which in the cannonball (CB) model are normal GRBs viewed farther off axis, the first peak-energy is shifted to the soft X-ray region while the second peak-energy moves to the MeV range. In far-off-axis GRBs, such as GRBs 980425 and 031203, the second peak may have been confused with the normal GRB peak. In most GRBs, which have been observed so far, the γ ray detectors ran out of statistics far below the second peak. However, in bright GRBs, the two peaks may be resolved by simultaneous measurements with SWIFT and GLAST.

1. Introduction

There is mounting evidence that long duration γ -ray bursts (GRBs) and X-ray flashes (XRFs) are both produced by highly relativistic and narrowly collimated jets ejected in supernova (SN) explosions akin to SN1998bw (see, e.g. Dar 2004 and references therein). Low-luminosity GRBs and XRFs seem to be ordinary GRBs whose γ -rays are much softer and their luminosity is much lower because they are viewed from angles relative to the jet which are a few times larger than the viewing angles of ordinary GRBs (see, e.g. Dar & De Rújula 2000,2003; Dado, Dar & De Rújula 2002,2004 and references therein). However, recently, the European Space Agency satellite INTEGRAL discovered GRB 031203 (Götz et al. 2004) at a redshift $z = 0.1055$ (Prochaska et al. 2004), the second nearest GRB after GRB 980425 at $z = 0.0085$ (Galama et al. 1998), which like GRB 980425 had extremely

¹dado@phep3.technion.ac.il, arnon@physics.technion.ac.il, dar@cern.ch.
Physics Department and Space Research Institute, Technion, Haifa 32000, Israel

low luminosity, but, contrary to the expectation, its γ -rays were not much softer than those of ordinary GRBs (Sazonov et al. 2004). GRB 031203 was produced in an SN very similar to SN1998bw which produced GRB 980425 (e.g. Malesani et al. 2004 and references therein). Consequently, the INTEGRAL observations were interpreted as evidence of a different class of GRBs, which are intrinsically very faint and include GRB 980425 and GRB 031203 (Sazonov et al. 2004; Soderberg et al. 2004; Woosley et al. 2004) and are produced in SNe akin to SN1998bw.

The spectrum of the soft γ -rays emitted in GRB 031203, which was measured with INTEGRAL, is well described by $dn_\gamma/dE \sim E^{-1.63 \pm 0.06}$ between 20 and 400 keV (Sazonov et al. 2004). But, its soft X-ray fluence in the 0.2-10 keV range, $F_x \sim 2.6 \pm 1.3 \times 10^{-7} \text{ erg s}^{-1}$, which was inferred (Watson et al. 2004) from its X-ray dust-scattered echo measured by XMM-Newton (Vaughan et al. 2004), is far above the INTEGRAL spectrum extrapolated to the soft X-ray region. It has been suggested that the soft X-ray fluence may have been overestimated (Prochaska et al. 2004; Sazonov et al. 2004).

However, in this letter we suggest an alternative interpretation of the INTEGRAL and XMM-Newton observations, which is based on the cannonball (CB) model of GRBs and XRFs (e.g. Dar & De Rújula 2003; Dado et al. 2004) and which removes the conflicts between the off-axis interpretation of the unusually low luminosity of GRB 031203 and its high peak-energy, and between its X-ray fluence extrapolated from the INTEGRAL measurements (Sazonov et al. 2004) and that inferred from the XMM-Newton measurements (Vaughan et al. 2004). We also demonstrate that this interpretation can explain other puzzling GRB observations.

The CB model assumes that in core-collapse SN, after the collapse of the stellar core into a neutron star or a black hole, an accretion disk or a torus is produced around the newly formed compact object, either by stellar material originally close to the surface of the imploding core and left behind by the explosion-generating outgoing shock, or by more distant stellar matter falling back after its passage (De Rújula 1987). A highly relativistic plasmoid (CB) of ordinary matter is emitted along the rotation axis when part of the accretion disk falls abruptly onto the compact object, as observed in microquasars (e.g. Mirabel & Rodriguez 1999; Rodriguez & Mirabel 1999 and references therein). A CB contains a thermal plasma with a power-law tail of knocked-on electrons from collisions with particles of the interstellar medium (ISM) and swept-in ISM electrons, which are Fermi-accelerated and cool quickly by synchrotron emission from the strong equipartition magnetic field in the CB to a steady state distribution, $dn_e/dE \sim E^{-(p+1)}$ with $p \sim 2.2$, in the CB rest frame. As the jet of CBs coasts through the “ambient light” permeating the surroundings of the parent SN, the electrons enclosed in the CBs Compton up-scatter photons to energies which, close to the

CBs direction of motion, correspond to the γ -rays of an ordinary GRB and less close to it, to the X-rays of an XRF (Dar & De Rújula 2003).

A CB also produces a narrow conical jet of high energy cosmic-ray electrons (and nuclei) along its motion (e.g., Dar & De Rújula 2003). These electrons are of two origins: swept-in ISM electrons which were Fermi-accelerated in the CB and escaped out into the ISM and ISM electrons which were scattered elastically by the CB. These cosmic-ray electrons produce a second peak in the spectral energy flux of GRBs and XRFs at a much higher energy, by inverse Compton scattering (ICS) of “ambient light” permeating the surroundings of the parent SN.

Thus, the internal CB electrons and the external cosmic ray electrons produce, by ICS of ambient light, a spectral-energy-density ($E^2 dn/dE$) of γ -rays with two peaks, like that observed in blazars (e.g., Padovani & Giommi 1995; Wehrle et al. 1998). Normal GRBs have their first peak-energy usually around a fraction of an MeV and then their second peak-energy at a much higher energy. Gamma-ray detectors on board satellites usually ran out of sensitivity/statistics well below the second peak-energy. However, high-energy photons with a flux much larger than that expected from the extrapolated decline of the first peak, have been discovered in a few cases of very bright GRBs with instruments on board CGRO (e.g. Hurley et al. 1994; Dingus et al. 1995; Gonzalez et al. 2003). In this letter we demonstrate that such a high energy component is well described by the CB model. Moreover, in very low-luminosity GRBs/XRFs, which are far off-axis GRBs, the first peak is in the keV range while the second peak is in the MeV range, i.e., the spacing between the peaks is relatively small and both peaks, or a significant fraction of both, may fall within the detection range of the ordinary XRF and GRB detectors. A score of low-luminosity XRFs were detected by CGRO, BeppoSAX and HETE II. But, in the case of HETE II, its FREGATE detector could go only to 400 keV. BATSE on board CGRO and GRBM on board BeppoSAX usually ran out of statistics/sensitivity well below the second peak, although a significant emission in excess of that expected from a simple power-law extrapolation of the low energy spectrum to higher energies was detected in some XRFs (see e.g., Frontera et al. 2004a,b). In this letter we first demonstrate that the peak-energy and the isotropic radiation energy emitted in the first peak satisfy well the simple correlation $(1+z) E_p \propto [E_\gamma^{iso}]^{1/3}$, predicted by the CB model. GRBs 980425 and 031203, which appear to be exceptions, could have been far-off-axis GRBs with a relatively small spacing between their keV and MeV peaks, and therefore appeared as GRBs with a “normal” peak-energy. We predict that simultaneous measurements with SWIFT and GLAST of the spectral energy flux of ordinary GRBs will resolve it into two peaks which shift to smaller energies during the GRB pulse.

2. The first peak

Let γ be the Lorentz factor of a CB and $\delta = 1/\gamma(1 - \beta \cos \theta)$ be its Doppler factor when viewed from an angle θ relative to its motion. In the CB model (Dar & De Rújula 2003), the observed peak-energy of γ -rays produced at a redshift z by ICS of thin thermal bremsstrahlung light around an SN, $dn_\gamma/dE \sim \exp(-E/T)/E$ with a typical energy $\epsilon_\gamma \sim T \sim 1 \text{ eV}$, is given by,

$$E_p \approx \frac{\gamma \delta}{1+z} T. \quad (1)$$

Under the assumption of isotropic emission in the CB rest frame, Doppler boosting and relativistic beaming yield a γ -ray fluence F_γ of a GRB pulse, which is proportional to δ^3 ,

$$F_\gamma \approx \frac{\delta^3 (1+z) E'_\gamma}{4\pi D_L^2}, \quad (2)$$

where E'_γ is the total γ -ray energy emitted in the CB rest frame. Consequently, E_γ^{iso} , the inferred ‘GRB isotropic γ -ray energy’ in a GRB pulse, **under the assumption of isotropic emission in the observer frame**, is

$$E_\gamma^{iso} = \frac{4\pi D_L^2}{1+z} F_\gamma \approx \delta^3 E'_\gamma. \quad (3)$$

In the CB model, the predicted GRB spectrum from ICS of ambient light with a thin thermal bremsstrahlung spectrum, by the electrons inside the CB, is,

$$\begin{aligned} \frac{dN_\gamma[1]}{dE} &\propto \left(\frac{T_{eff}}{E}\right)^\alpha e^{-E/T_{eff}} + b(1 - e^{-E/T_{eff}}) \left(\frac{T_{eff}}{E}\right)^\beta \\ \alpha &\approx 1; \quad \beta = \frac{p+2}{2} \approx 2.1, \end{aligned} \quad (4)$$

where $T_{eff} = \gamma \delta T/(1+z)$ and b is a dimensionless constant. The values of α and β may deviate from their indicated values, because the ambient radiation may deviate from a thin thermal bremsstrahlung, and the power-law index of the accelerated and knocked-on electrons after cooling may be larger than $p+1 = 3.2$ and increase with time. Also the temperature of the ambient light seen by the CB is decreasing with distance, approximately as

$$T_{eff}(t) \sim T_{eff}(0) \{1 - \exp[-(t_0/t)^2]\}, \quad (5)$$

where t_0 is a constant. As was shown in Dar & De Rújula 2003 and in Dado et al. 2004, Eq. (4) is practically indistinguishable from the phenomenological Band function (Band et al. 1993) and it is in good agreement with the measured spectrum of the photons in the first peak of the spectral-energy-density of ordinary GRBs and XRFs.

If core collapse SNe and their environments were all identical, and if their ejected CBs were also universal in number, mass, Lorentz factor and velocity of expansion, all differences between GRBs would depend only on the observer’s position, determined by z and the angle of observation, θ . For a distribution of Lorentz factors that is narrowly peaked around $\gamma \simeq 10^3$, the θ -dependence is in practice the dependence on δ , the Doppler factor. Hence Eqs. (1),(3) yield the correlation (Dar and De Rújula 2000; 2003),

$$(1+z) E_p \propto [E_\gamma^{iso}]^{1/3}. \quad (6)$$

In Fig. 1 we plot $(1+z) E_p$ as function of E_γ^{iso} for GRBs/XRFs of known redshift, as compiled by Amati (2004) and by Ghirlanda et al. (2004), and the best fitted power-law, $(1+z) E_p = a [E_\gamma^{iso}]^\alpha$. The best fitted power-law index obtained with the CERN standard program MINUIT, $\alpha = 0.335 \pm 0.06$ and $\alpha = 0.345 \pm 0.07$, for the Amati (2004) and Ghirlanda et al. (2004) compilations, respectively, is in agreement with Eq. (6) but not with the original relation, $(1+z) E_p \sim a [E_\gamma^{iso}]^{0.52 \pm 0.06}$ proposed by Amati et al. (2002), which, as far as we know, has no theoretical derivation. We also plot in Fig. 1 parallel lines between which the observed values are expected to be spread because of the spread in the ‘standard candle’ properties. The correlation predicted by the CB model is well satisfied except for GRB 980425 and GRB 031203.

3. The second peak

In the CB model, the ISM in front of the CBs is ionized by the beamed radiation from the highly relativistic CBs. The turbulent magnetic fields in the CBs accelerate the ionized ISM particles, which they gather on their path, to an initial distribution, $dn_e/dE \sim E^{-p}$ with $p \sim 2.2$ in the CB rest frame. In a steady state situation, the electrons, which are trapped in the CB by its internal magnetic fields and cool quickly by synchrotron emission, reach a distribution, $dn_e/dE \sim E^{-(p+1)}$, while the electrons which escape the CB must have the hard “injection spectrum”, $dn_e/dE \sim E^{-p}$. Their cooling time in the ISM is much longer than that of the electrons which are trapped magnetically inside the CBs, because the ISM magnetic field is smaller by many orders of magnitude than that inside the CBs.

In the CB rest frame, the Lorentz boosted ambient light undergoes Compton scattering from these two distributions and produces the first peak with the “Band function” shape and a hard tail at higher energies with practically a “time-independent” spectral index 1.6,

$$\frac{dn_\gamma}{dE} \propto E^{-(p+1)/2} \sim E^{-1.6}. \quad (7)$$

The ISM electrons which are scattered elastically by the highly relativistic CBs in the

direction of the observer have an approximate lab energy, $E_e \sim \gamma \delta m_e$. Because of their very large Lorentz factor, $\Gamma_e \sim \gamma \delta$, the ambient photons which they scatter have much higher energies than those of the photons scattered by “cold” electrons in the CBs, and they are narrowly beamed along the electrons’ direction of motion. Hence, in the Thomson regime they have approximately the thin thermal bremsstrahlung distribution of the ambient light boosted by $\sim 4 \gamma^2 \delta^2 / 3$:

$$\frac{E^2 dN_\gamma}{dE} \propto E e^{-E/T_{eff}}, \quad (8)$$

where $T_{eff} = (4/3) \gamma^2 \delta^2 T / (1+z)$. For $E \ll T_{eff}$, the low energy side of the spectral energy density of this “second peak” has the simple shape, $E^2 dN_\gamma / dE \propto E$.

During the GRB phase, the radius of a CB increases linearly with the distance x from the explosion site while the density of the ambient light decreases like $1/x^2$. Because of the slow cooling rate in the ISM, the accumulated number of high energy electrons in the narrow beam produced by the highly relativistic CB is proportional to the ISM mass swept up by the CB. Thus, for a constant density-profile, the magnitude of the spectral energy density, which is proportional to the product of the swept-up mass and the density of the ambient light, increases linearly with distance, whereas it decreases like $1/x$ for a density-profile of a stellar wind which blows constantly and produces a density $n_e \propto 1/x^2$. During the GRB the deceleration of the CBs is negligible and the observer time is proportional to the distance. Thus, for a constant density-profile the spectral energy density increases with time like t , whereas it decreases like $1/t$ for a “windy” density-profile.

In the Thomson regime, the peak-energies are given approximately by $E_p \approx T_{eff} / (1+z)$. Consequently, $E_p[1]$ and $E_p[2]$, the peak energies of the first and second peak, respectively, are related through

$$E_p[2] \approx \frac{(1+z) [E_p[1]]^2}{T}. \quad (9)$$

For ordinary GRBs with $E_p[1] \sim 200 \text{ keV}$ and $z \sim 1$, Eq. (9) yields $E_p[2] \sim 100 \text{ GeV}$, whereas for a very dim XRF with $E_p[1] \sim 1 \text{ keV}$ and $z \ll 1$, it yields, $E_p[2] \sim 1 \text{ MeV}$. During each GRB pulse, both peak-energies, being proportional to $T(t)$, decrease during a GRB pulse like, $T_{eff}(0) [1 - \exp[-(t_0/t)^2]]$.

4. Comparison with observations

There are very few cases where GRBs were measured over a very broad energy range and allow a comparison between the CB model predictions and the observations. Here we shall limit the comparison to two GRBs of particular interest.

4.1. GRB 941017

Gonzalez et al. (2003) reported the discovery a high energy spectral component in GRB 940117 with an energy flux density, $E^2 dn/dE \sim E$, peak energy $E_p[2] \geq 200 \text{ MeV}$ and a fluence ≥ 3 times that of the the first normal GRB component. Their results are shown in Fig. 2 borrowed from Gonzalez et al. 2003. The theoretical lines are best fitted Band functions plus a power-law contribution (Gonzalez et al. 2003). But, these lines are indistinguishable from the bes-fitted CB model double-peak spectral-energy-density which is a sum of $E^2 dn/dE$ with dn/dE as given by Eq. (4) and of a second component which is described by Eq. (8) with $E \ll T_{eff}[2]$.

4.2. GRB 031203

GRB 031203 (Götz et al. 2003) was produced in an SN explosion similar to SN1998bw (Bersier et al. 2004; Cobb et al. 2004; Malesani et al. 2004; Thomsen et al. 2004; Gal-Yam et al. 2004) and had an unusually low inferred isotropic luminosity. Its soft X-ray emission, which was inferred from modeling its measured dust scattered echo by XMM-Newton, yielded (Watson et al. 2004; Vaughan et al. 2004)), $F_x \sim 2.6 \pm 1.3 \times 10^{-6} \text{ erg cm}^{-1}$, in the 0.7-5 keV range, far above the extrapolation of the INTEGRAL spectrum. This can be seen from the upper panel of Fig. 3 where we demonstrate a CB model fit to the spectral energy density of GRB 031203 which consists of two terms, $E^2 dn/dE$ with dn/dE as given by Eq. (4) for a low-energy peak with $E_p \approx T_{eff} \approx 3 \text{ keV}$ and $b = 10^{-2}$, and a hard-tail contribution as given by Eq. (7). The normalization of both components were adjusted to fit the observational data. In the lower panel of Fig. 3b we compare the photon spectral index of the hard tail component, $(p+1)/2 \approx 1.6$, as predicted by the CB model, Eq. (7), and its measured value by INTEGRAL (Sazonov et al. 2004).

5. Conclusions

In the CB model, the internal population of electrons produce the ordinary peak in the spectral-energy-density of GRBs/XRFs and the external population of high energy (cosmic ray) electrons, which are accelerated by the CBs, produce a second GRB peak at a much higher energy through inverse Compton scattering of ambient light permeating the surroundings of the parent SN. We have shown that for GRBs with known redshift, usually the peak-energy and the isotropic radiation energy of the first, ordinary peak satisfy well the simple correlation $(1+z) E_p \propto [E_\gamma^{iso}]^{1/3}$, predicted by the CB model. However, GRBs

980425 and 031203 have peak-energies much larger than those expected in the CB model from their very low luminosities. But, in very low luminosity GRBs and XRFs which, in the CB model, are ordinary GRBs viewed far off axis, the second peak moves down to the MeV region and can dominate the soft γ -ray region. We have shown that GRBs 031203 and GRB 980425 could have been such far-off-axis GRBs with their first peak in the soft X-ray region but with a contribution from their second broad peak which dominates their X-ray and γ -ray emission, as shown here explicitly for GRB 031203. We have also shown that the high energy component which was discovered in GRB 941017 is well explained by the second peak. Finally, we predict that simultaneous measurements with SWIFT and GLAST will resolve the spectral energy flux of ordinary GRBs/XRFs into a double peak spectrum whose peaks move to lower energies during the GRB pulse.

Acknowledgement: Useful comments by Avishai Gal-Yam, Kevin Hurley and Simon Vaughan are gratefully acknowledged. This research was supported in part by the Asher Fund for Space Research at the Technion.

REFERENCES

- Amati, L., et al. 2002, A&A, 390, 81
- Amati, L. 2004, astro-ph/0405318
- Band, D., et al. 1993, ApJ, 413, 281
- Bersier, D., et al. 2004, GCN Circ. 2544
- Cobb, B. E., et al. 2004, astro-ph/0403510
- Dado, S., Dar A. & De Rújula, A. 2002, A&A, 388, 1079
- Dado S., Dar A. & De Rújula A. 2004, A&A, 422, 381
- Dar, A. 2004, astro-ph/0405386
- Dar, A. & De Rújula, A. 2000, astro-ph/0012227
- Dar, A. & De Rújula, A. 2003, astro-ph/0308248 (Phys. Rep. in press)
- De Rújula, A. 1987, Phys. Lett. 193, 514
- Dingus, B. L., et al. 1995, Ap&SS, 231, 187
- Frontera, F., et al. 2004b, astro-ph/astro-ph/0406575

- Frontera, F., et al. 2004, astro-ph/0408436
- Galama, T. J., et al. 1998, Nature, 395, 670
- Gal-Yam, A., et al. 2004, ApJ, 609, L59
- Gonzalez, M. M., et al. 2003, Nature, 424, 749,
- Götz, D., et al. 2003, GCN 2459
- Ghirlanda, G., Ghisellini, G. & Lazzati, D, 2004, astro-ph/0405602
- Hurley, K., et al. 1994, Nature, 372, 652
- Malesani, D., et al. 2004, astro-ph/0405449
- Padovani, P. & Giommi P. 1995, ApJ, 444, 567
- Prochaska, J. X., et al. 2004, ApJ, 611, 200
- Sazonov, S. U., Lutovinov, A. A., & Sunyaev, R. A. Nature, 2004, 430, 646
- Soderberg, A. M., et al. 2004, Nature, 430, 648
- Thomsen, B., et al. 2004, astro-ph/0403451
- Vaughan, S., et al. 2004, ApJ, 603, L5
- Watson, D., et al. 2004, ApJ, 605, L101
- Wehrle, et al. 1998, ApJ, 497, 178
- Woosley, S., 2004, Nature, 430, 623
- Yamazaki, R., Yonetoku, D. & Nakamura, T. 2003, ApJ, 594, L79




Fig. 1.— The observed peak-energy corrected for redshift, $(1+z)E_p$ as function of the inferred isotropic radiation energy emitted at redshift z for GRBs/XRFs of known redshift as compiled by (a) Amati (2004) and by (b) Ghirlanda et al. (2004). The thick lines are the best fitted power-laws: $A E^{0.334 \pm 0.06}$ and $A E^{0.345 \pm 0.07}$, respectively. The dotted lines border the estimated spread (a factor of ~ 4) in the isotropic radiation energy due to the spread in the number of CBs and their physical properties in GRBs. The point due to GRB 980425 (as inferred by Yamazaki et al. 2003) is far above the theoretical expectation.




Fig. 2.— The spectral energy density of GRB 941017 obtained by Gonzalez et al. (2003) with BATSE-LAD (crosses) and EGRET-TASC (filled circles) on board CGRO. The CB model fit for the first peak is indistinguishable from the “Band function” fit (dashed line) by Gonzalez et al. (2003). The rising side of the second peak is well described by the CB model prediction as given by Eq. (8). The fits are shown for the time intervals (-18)-14s (a);




Fig. 3.— Top: Comparison between the observed spectral energy density of GRB 031203 and a CB model fit given by Eq. (4) with $E_p \approx T_{eff} \approx 3 \text{ keV}$ and $b = 10^{-2}$, and a “hard tail” as given by Eq. (7). The X-ray fluence is that inferred by Vaughan et al. (2004) from the dust echo observed with XMM-Newton. The INTEGRAL measurements are those reported by Sazonov et al. (2004). Bottom: Comparison between the the photon spectral index as obtained from single power-law fit to the photon spectrum measured with INTEGRAL (Sazonov et al. 2004) and the CB model prediction for the hard-tail component in GRBs generated by cosmic ray electrons which were accelerated in the CBs.